

Oscillating Chemical Reactions Lab

Math 230

due Friday, December 8 at 4pm

Certain chemical reactions oscillate rather than approach equilibrium. In this lab you will investigate a system of differential equations that models such a reaction. The system depends on a parameter, B , and you are to describe qualitatively what happens when B passes through a certain threshold value. There are two parts to this lab. After making the appropriate computations in Part I, use technology to view the phase plane for the system and report on the qualitative behavior of the system in Part II.

Suppose that iodide and chlorine dioxide are each poured into a reaction vessel at a certain rate and are mixed thoroughly, causing a reaction to occur. The excess solution is removed so that the amount of reactant in the vessel remains constant. Let x be the amount of iodide and y the amount of chlorine dioxide in the vessel at time t . A mathematical model for a chlorine dioxide-iodide open system reaction is provided by the following system of (nonlinear) differential equations:

$$\begin{aligned}\frac{dx}{dt} &= -x + 10 - \frac{4xy}{1+x^2} \\ \frac{dy}{dt} &= B \left(x - \frac{xy}{1+x^2} \right)\end{aligned}$$

The quantity B is a parameter that we will let vary between 1 and 6.

Part I: Calculations

Do the following calculations first on scratch paper, and then write your solutions clearly and neatly.

1. Find all of the equilibrium points for this system for all values of the parameter B .
Hint: You should find exactly one equilibrium point for each value of B , and this equilibrium point should not depend on B . If this is not the case, check your work carefully!
2. Compute the Jacobian matrix of the system at a general point (x, y) . *Be careful here:* You must have the correct Jacobian matrix in order to proceed with the following questions.
3. Now compute the Jacobian matrix at the equilibrium point you found in 1. This matrix should depend on B . Then find the eigenvalues of this matrix. These should also depend on B .
4. Classify these equilibria as spiral sinks, real sources, saddles, or whatever. Your answer should include a range of B values for each type that occurs. Remember, we are assuming that $1 \leq B \leq 6$.
5. Next, answer the following question: At what value of B does a bifurcation occur, and what happens at this bifurcation?
6. Sketch the curve in the trace-determinant plane given by this matrix as B varies.
7. Sketch the nullclines for this system and indicate the direction of the vector field in the regions between these nullclines. How do these nullclines depend on the parameter B ? It suffices to restrict attention here to the region $0 < x < 5$, $3 < y < 8$.

Part II: Observations

Use technology to help you answer the following questions. For example, you could use *Mathematica* to numerically solve the system with various initial conditions and display solution curves in the phase plane. See sample *Mathematica* code in the reference section below. Explain your answers clearly.

8. Discuss the behavior of solutions of the system for the parameter value $B = 6$. Answer the question: What happens to the chemicals in the vessel as time passes?

9. Discuss the behavior of solutions of the system for the parameter value $B = 1$. Answer the question: What happens to the chemicals in the vessel as time passes?
10. As the parameter B varies between 1 and 6, we know that there is a bifurcation (a qualitative change in the solutions) at some B value, and from the linearization in Part I, we know what to expect there. Write a few sentences explaining what this bifurcation entails, both mathematically and chemically. That is, as the parameter B passes through the bifurcation value, explain what happens in the phase plane and what this means in terms of the chemicals involved in the reaction.

Reference: Phase Plots in *Mathematica*

The following code will help you plot the phase plane, with vector streams and a sample solution on the same diagram.

- Create a stream plot and save it as `splot` (assume B already has a value between 1 and 6):

```
splot = StreamPlot[{-x + 10 - 4x*y/(1 + x^2), B(x - x*y/(1 + x^2))},
  {x, 0, 5}, {y, 3, 8}]
```

- Numerically solve the system with initial condition $x(0) = r$ and $y(0) = s$; save the solution as `sol`:

```
eqns = {x'[t] == -x[t] + 10 - 4 x[t] y[t]/(1 + (x[t])^2),
  y'[t] == B (x[t] - x[t] y[t]/(1 + (x[t])^2)),
  x[0] == r, y[0] == s};
sol = NDSolve[eqns, {x, y}, {t, 0, 20}]
```

- Create a parametric plot of the solution; save it as `pplot`:

```
pplot = ParametricPlot[Evaluate[{x[t], y[t]} /. sol], {t, 0, 20},
  PlotStyle -> Red]
```

- Plot the stream plot and parametric plot together:

```
Show[splot, pplot]
```

Lab Report

Type your answers to the numbered items above in a document. Make sure that you clearly state the solution to each numbered item! You may use L^AT_EX or a word processor. You may draw diagrams by hand. *Do not* turn in a Mathematica file. If you submit your report electronically, please submit a PDF file.

Grading

Your lab report will be graded out of 40 points, based on the following criteria:

1. **Correctness:** Answers and supporting work are mathematically accurate. In particular, the bifurcations have been correctly identified. (20 points)
2. **Completeness:** Assigned questions are answered completely, with appropriate details and justification to support the answers. (8 points)
3. **Clarity:** Explanations are clear and concise. English sentences are used along with equations to explain mathematical reasoning. (6 points)
4. **Presentation:** Work is presented in a typed document that is neat, organized, and easy to read. (6 points)