

1. Let  $X \sim \text{Geometric}(p)$ . Compute the mgf  $M_X(t)$ . Try to write the mgf without a summation.

(a) What is the mgf  $M_X(t)$ ? Write your answer as a sum.

$$M_X(t) = E(e^{tX}) = \sum_{k=1}^{\infty} e^{tk} \underbrace{(1-p)^{k-1} p}_{P(X=k)}$$

(b) Your summation in part (a) should be a geometric series. Use the geometric sum formula to write the mgf  $M_X(t)$  without a sum.

Geometric series:  $\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r}$

$$M_X(t) = \sum_{k=1}^{\infty} p e^{tk} (1-p)^{k-1} = \underbrace{pe^t}_{\text{first term } a = pe^t} + pe^{2t}(1-p) + pe^{3t}(1-p) + pe^{4t}(1-p) + \dots$$

Common ratio:  $r = e^t(1-p)$

Thus,  $M_X(t) = \frac{pe^t}{1 - e^t(1-p)}$

Alternative approach:

$$M_X(t) = \frac{p}{1-p} \sum_{k=1}^{\infty} e^{tk} (1-p)^k = \frac{p}{1-p} \sum_{k=1}^{\infty} \underbrace{(e^t(1-p))^k}_{\text{geometric series: } a=r=e^t(1-p)}$$

$$= \frac{p}{\cancel{1-p}} \cdot \frac{\cancel{e^t(1-p)}}{1 - e^t(1-p)} = \frac{pe^t}{1 - e^t(1-p)}$$

2. Suppose random variable  $X$  has probability mass function  $P(X = x) = \frac{27}{40} \left(\frac{1}{3}\right)^x$ , for integers  $0 \leq x \leq 3$ .

(a) Verify that this is a valid probability mass function.

• nonnegative probabilities

• sum is:

$$\frac{27}{40} \left( 1 + \frac{1}{3} + \frac{1}{9} + \frac{1}{27} \right) = \frac{27}{40} \left( \frac{1 - \left(\frac{1}{3}\right)^4}{1 - \frac{1}{3}} \right) = \frac{27}{40} \left( \frac{\frac{80}{81}}{\frac{2}{3}} \right) = \frac{27}{40} \cdot \frac{40}{27} = 1$$

4 terms of a geometric series with  $r = \frac{1}{3}$

$a + ar + ar^2 + \dots + ar^{n-1} = a \frac{1-r^n}{1-r}$

(b) Find the moment generating function  $M_X(t)$ . Try to write the mgf without a summation.

$$M_X(t) = E(e^{tX}) = e^0 \cdot \frac{27}{40} + e^t \cdot \frac{9}{40} + e^{2t} \cdot \frac{3}{40} + e^{3t} \cdot \frac{1}{40} = \frac{27}{40} \cdot \frac{1 - (\frac{e^t}{3})^4}{1 - (\frac{e^t}{3})} = \frac{81 - e^{4t}}{40(3 - e^t)}$$

common ratio:  $r = \frac{e^t}{3}$

(c) Compute  $M'_X(0)$ . Does your answer agree with the expected value of  $X$  computed directly from the pmf?

$$M_X(t) = \frac{81 - e^{4t}}{40(3 - e^t)} \quad M'_X(t) = \frac{40(3 - e^t)(-4e^{4t}) - (81 - e^{4t})(40)(-e^t)}{40^2(3 - e^t)^2} \quad \text{Use technology!}$$

$$M'_X(0) = \frac{40(2)(-4) - (80)(40)(-1)}{40^2(2)^2} = \frac{72}{160} = \frac{9}{20}$$

$$E(X) = 0 + 1 \cdot \frac{9}{40} + 2 \cdot \frac{3}{40} + 3 \cdot \frac{1}{40} = \frac{9 + 6 + 3}{40} = \frac{9}{20} \quad \leftarrow \text{same!}$$

↙ This problem is in the next class video.

3. Let  $X$  be a discrete random variable with mgf  $M_X(t)$ , and let  $Y = aX + b$ . Write the definition of  $M_Y(t)$ . Replace  $Y$  with  $aX + b$  and simplify the expected value to show that  $M_Y(t) = e^{bt} M_X(at)$ .

$$M_Y(t) = E(e^{tY}) = E(e^{t(aX+b)}) = E(e^{atX+bt}) = e^{bt} E(e^{atX}) = e^{bt} M_X(at)$$

If  $Y = aX + b$ , then  $M_Y(t) = e^{bt} M_X(at)$ .