


1. Let  $X$  be a continuous random variable with pdf


$$f(x) = \begin{cases} x & 0 \leq x \leq 1, \\ 2-x & 1 \leq x \leq 2, \\ 0 & \text{otherwise.} \end{cases}$$

(a) Sketch the pdf of  $X$ .



(b) Without computing anything, sketch cdf of  $X$ .


The cdf must look something like this:




(c) What is  $P(X < 1.5)$ ?

Find the area under the pdf left of  $x = 1.5$ .

$$P(X < 1.5) = \int_0^{1.5} f(x) dx = \frac{7}{8}$$



(d) Find a value  $\eta_{0.75}$  such that  $P(X \leq \eta_{0.75}) = 0.75$ .



Find  $c$  such that  $c$  has area  $\frac{1}{4}$ .

$$\begin{aligned} \frac{c^2}{2} &= \frac{1}{4} \\ c^2 &= \frac{1}{2} \\ c &= \frac{\sqrt{2}}{2} \end{aligned}$$

$$\begin{aligned} \text{so } \eta_{0.75} &= 2 - \frac{\sqrt{2}}{2} \\ &\approx 1.293 \end{aligned}$$

(e) Give a formula for the cdf of  $X$ .


If  $0 \leq x \leq 1$ , then  $F(x) = \int_0^x y dy = \frac{1}{2} y^2 \Big|_{y=0}^{y=x} = \frac{1}{2} x^2$   
cdf:  $P(X \leq x)$       pdf

If  $1 \leq x \leq 2$ , then  $F(x) = \frac{1}{2} + \int_1^x (2-y) dy = \frac{1}{2} + [2y - \frac{1}{2}y^2]_{y=1}^{y=x} = 2x - \frac{x^2}{2} - 1$   
 $P(X \leq 1)$        $P(1 \leq X \leq x)$

Thus: 
$$F(x) = \begin{cases} 0 & \text{if } x < 0 \\ \frac{1}{2}x^2 & \text{if } 0 \leq x \leq 1 \\ 2x - \frac{x^2}{2} - 1 & \text{if } 1 < x \leq 2 \\ 1 & \text{if } 2 < x \end{cases}$$

2. Suppose that a continuous random variable  $X$  has pdf  $f(x) = kx(4-x)$  for  $0 \leq x \leq 4$ , and  $f(x) = 0$  otherwise.

(a) Sketch the pdf of  $X$ . Then, without computing anything, sketch the cdf of  $X$ .



(b) What is the value of  $k$ ?

Remember that the pdf must integrate to 1:

$$\int_0^4 kx(4-x) dx = \frac{32}{3}k = 1, \quad \text{so } k = \frac{3}{32}.$$

(c) Find  $P(X < 1 \text{ or } X > 3)$ .


$$\begin{aligned} P(X < 1 \text{ or } X > 3) &= \int_0^1 f(x) dx + \int_3^4 f(x) dx = 2 \int_0^1 \frac{3}{32} x(4-x) dx \\ &\quad \text{by symmetry of } f(x) \text{ about } x=2 \\ &= \frac{3}{16} \left[ 2x^2 - \frac{x^3}{3} \right]_0^1 = \frac{3}{16} \left( 2 - \frac{1}{3} \right) = \frac{5}{16} \end{aligned}$$

3. Suppose that the cdf of a random variable  $X$  is  $F(x) = 1 - e^{-5x}$  for  $x > 0$ , and  $F(x) = 0$  otherwise.

(a) What is the pdf of  $X$ ? Sketch both the pdf and the cdf.

Differentiate the cdf to find the pdf:

$$f(x) = \frac{d}{dx} F(x) = \frac{d}{dx} (1 - e^{-5x}) = 5e^{-5x} \quad \text{for } x > 0.$$



(b) What is  $P\left(\frac{1}{4} < X < \frac{1}{3}\right)$ ? Can you get this from either the cdf or the pdf?

$$\text{cdf: } P\left(\frac{1}{4} < X < \frac{1}{3}\right) = F\left(\frac{1}{3}\right) - F\left(\frac{1}{4}\right) = (1 - e^{-5/3}) - (1 - e^{-5/4}) = e^{-5/4} - e^{-5/3} \approx 0.098$$

$$\text{pdf: } P\left(\frac{1}{4} < X < \frac{1}{3}\right) = \int_{1/4}^{1/3} f(x) dx = \text{same}$$

4. Random variable  $X$  has pdf  $f(x) = \begin{cases} ax + bx^2 & 0 < x < 1, \\ 0 & \text{otherwise.} \end{cases}$

Furthermore,  $P\left(X < \frac{1}{2}\right) = \frac{3}{16}$ . What is the median of  $X$ ?

We need:  $\int_0^1 (ax + bx^2) dx = \frac{a}{2} + \frac{b}{3} = 1 \Rightarrow 3a + 2b = 6$

$$\int_0^{\frac{1}{2}} (ax + bx^2) dx = \frac{a}{2} \cdot \frac{1}{4} + \frac{b}{3} \cdot \frac{1}{8} = \frac{a}{8} + \frac{b}{24} = \frac{3}{16} \Rightarrow 3a + b = \frac{9}{2}$$

$$b = \frac{3}{2}, a = 1$$

Median:  $\int_0^{\eta} (x + \frac{3}{2}x^2) dx = \frac{1}{2}\eta^2 + \frac{1}{2}\eta^3 = \frac{1}{2} \Rightarrow \eta^2 + \eta^3 = 1$

So we need  $\eta^3 + \eta^2 - 1 = 0$ , or  $\eta \approx 0.755$

$$\text{Exact: } \eta = \frac{1}{3} \left[ -1 + \sqrt[3]{\frac{25}{3} - \frac{3\sqrt{69}}{2}} + \sqrt[3]{\frac{1}{2}(25 + 3\sqrt{69})} \right]$$