


1. Simulate 10,000 averages, each of k samples from a $\text{Unif}[0,1]$ distribution. Make a histogram of the 10,000 averages. Start with $k = 1$ and then try larger values of k . How does the shape of the histogram depend on k ?


larger $k \Rightarrow$ closer to normal



Histogram of 10,000 samples, each a sum of 5 $\text{Unif}[0,1]$ random variables.

2. Repeat the previous simulation, but now replace $\text{Unif}[0,1]$ with a different distribution of your choice. What is the shape of the histogram? How does it depend on k ?

again,
larger $k \Rightarrow$ closer to normal



Histogram of 10,000 samples, each a sum of 40 $\text{Hypergeometric}(n=8, M=20, N=50)$ random variables.

3. Let X_1, X_2, \dots, X_{300} be iid random variables with mean μ_X and standard deviation σ_X . Also let $T = X_1 + X_2 + \dots + X_{300}$ and $\bar{X} = \frac{T}{300}$.

(a) What are μ_T , σ_T , $\mu_{\bar{X}}$, and $\sigma_{\bar{X}}$?

$$\mu_T = 300\mu_X$$

$$\sigma_T = \sigma_X \sqrt{300}$$

$$\mu_{\bar{X}} = \mu_X$$

$$\sigma_{\bar{X}} = \frac{\sigma_X}{\sqrt{300}}$$

(b) What distributions are good approximations for T and \bar{X} ?

T is approx. $N(300\mu_x, \sigma_x\sqrt{300})$, \bar{X} is approx $N(\mu_x, \frac{\sigma_x}{\sqrt{300}})$

4. Use the Convolve function in Mathematica to plot the pdf of $X_1 + X_2 + \dots + X_n$, where each $X_i \sim Unif[0,1]$ and $n \in \{1, 2, 3, 4, 5, 6\}$. Compare each pdf with the pdf of a normal distribution.

See Mathematica notebook.

5. A farm packs tomatoes in crates. Individual tomatoes have mean weight of 10 ounces and standard deviation of 3 ounces. Estimate the probability that a crate of 40 tomatoes weighs between 380 and 410 ounces.

T_{40} is approximately $N(400, 18.97)$

$P(380 < T_{40} < 410) \approx 0.555$

R: $\text{pnorm}(410, 400, 18.97) - \text{pnorm}(380, 400, 18.97)$